(30) Priority Data:

N

ġ





INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 7:		(11) International Publication Number:	WO 00/68418
	A1	(43) International Publication Date:	6 November 2000 (16.11.00)

(21) International Application Number:	PCT/GB00/01740
(21) International Expenses	•

5 May 2000 (05.05.00) (22) International Filing Date:

GB 10 May 1999 (10.05.99) 9910811.0

(71) Applicant (for all designated States except US): MEDICAL RESEARCH COUNCIL [GB/GB]; 20 Park Crescent, London W1N 4AL (GB).

(72) Inventors; and (75) Inventors/Applicants (for US only): BRUNE, Martin, Hermann, Klemens [DE/GB]; National Institute for Medical Research, The Ridgeway, Mill Hill, London NW7 1AA (GB). CORRIE, John, Edgar, Thomas [GB/GB]; National Institute Ū for Medical Research, The Ridgeway, Mill Hill, London ũ NW7 1AA (GB). WEBB, Martin, Ronald [GB/GB]; Na-W tional Institute for Medical Research, The Ridgeway, Mill Hill, London NW7 1AA (GB).

Agents: HALLYBONE, Huw, George et al.; Carpmaels & Ransford, 43 Bloomsbury Square, London WC1A 2RA (GB).

(81) Designated States: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).


Published

With international search report.

Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.

BEST AVAILABLE COPY





(57) Abstract

Nucleoside diphosphates (NDPs) are detected or measured by following the dephosphorylation of the phosphoenzyme form of nucleoside diphosphate kinase (NDPK), and nucleoside triphosphates (NTPs) are detected or measured by following the phosphorylation of NDPK to its phosphoenozyme form. A typical process involves (a) causing NDP in sample to bind to NDPK phosphoenzyme, or causing NTP in sample to phosphorylate NDPK and (b) detecting a change in a characteristic of the enzyme which differs between its phosphorylated and unphosphorylated forms. This may be aided by labelling the enzyme. Quantitative data can be obtained. Both in vivo and in vitro measurements can be made.